

WHS REMOTE VETERINARY CARE DISCHARGE FORM

Your pet had the following procedures performed today:

Rabies vaccine

DH2PPV vaccine

Parasite treatment:

FCRVP vaccine

Suprelorin 9.4mg temporary sterilisation implant

Thank you for providing the proper care for your dog or cat!

Here are some tips on how to provide the best care for your furry friend:

It is normal to have some swelling and/or pain at then injections sites on either shoulder or between the shoulder blades for 1-2 weeks.

If you pet has any symptoms of being low energy, not eating, vomiting, profuse diarrhea, swelling of the face, difficulty breathing please call or text _____

If before (date). If you have questions or concerns regarding your pets health related to the procedures performed today after the above date, please call the Winnipeg Humane Society Clinic at 204-982-2036 or email clinic@winnipeghumanesociety.ca.

If your dog received a temporary sterilisation Suprelorin 9.4mg implant today, your pet needs to be kept away from dogs of the opposite sex for 8 weeks post implant and then will have temporary sterilisation for up to 12 months. Male and female dogs need to be kept separate for 8 weeks post implant as there is a delay in efficacy of the implant. After 8 weeks, the implant will provide temporary sterilization of your dog for up to 12 months in males and females. The implant may initially cause your female dog to go into heat so keeping her away from males is particularly important. If she happens to be bred in the 8 weeks after implantation, she will likely abort the litter before birth. This implant may predispose your female dog to an infection in the uterus called pyometra which would result in her needing to be surgically sterilized. Other potential complications in females include prolonged bloody vaginal discharge or heat. Swelling at the injection site for 14 days post implantation can be normal. The implant is only a temporary method of sterilization lasting up to 12 months. Hair loss may also be noted at the injection site. Dogs with known dog-dog /dog-person/dog-animal aggression may experience an increase in aggressive behaviour after implant sterilization.